


The Gomphidae of Algeria and the Maghreb: status, ecology and conservation (Insecta: Odonata)

Djamila Hamzaoui^a, Mouna Hafiane^a, Mohamed Mebarki^a, Abdeslem Arab^a,
Ahmed H. Alfarhan^b and Boudjéma Samraoui^{b,c,*} 

^aLaboratory of Dynamics and Biodiversity, FSB, USTHB, BP 32 El Alia, Bab Ezzouar, Algiers, Algeria;

^bCenter of Excellence for Research in Biodiversity, King Saud University, Riyadh, Saudi Arabia;

^cLaboratoire de Recherche et de Conservation des Zones Humides, University of Guelma, Guelma, Algeria

(Received 5 February 2015; accepted 18 April 2015)

A survey of the Gomphidae of Algeria and the Maghreb was carried out during the period 2013–2014. Sampling of eight main wadis across northern Algeria was undertaken and adults and exuviae were recorded. The survey yielded six species of Gomphidae. Among these, we report on the rediscovery of the Critically Endangered *Lindenia tetraphylla* in Algeria, recorded in the nineteenth century and deemed to have been extinct after an absence of more than a century and a half. An exuvia was collected at Wadi Saoura, which constitutes the first proof of the breeding of this species in Algeria and the third record for North Africa. We also recorded a pale form of *Onychogomphus uncatus*, morphologically distinct from typical Moroccan and European phenotypes, suggesting some degree of subspeciation and inviting further taxonomical investigations of the genus *Onychogomphus* in North Africa. Due to increasing water demand, Gomphidae and their habitats are under great pressure in the Maghreb. Specific threats and conservation measures are discussed.

Keywords: Algeria; anthropogenic impact; ecology; Gomphidae; Odonata; *Onychogomphus uncatus*; *Lindenia tetraphylla*; niche partitioning; North African wadi; vicariance

Introduction

Odonata have complex life cycles with discrete life stages spent in distinct habitats (Corbet, 1999). Dragonfly larvae are relatively large obligate predators which may have a considerable impact on community structure of freshwater ecosystems (Woodward & Hildrew, 2001, 2002). Among Odonata, gomphids are an important component of the biota of running waters and their numbers and diversity are considered a reliable indicator of a healthy fluvial system (Ferrerias-Romero, Márquez-Rodríguez, & Ruiz-García, 2009). The need to update our knowledge of the Gomphidae of Algeria is clear as some species have not been recorded since the nineteenth or early twentieth centuries. Their recent assessment by IUCN was a “best of a bad job” undertaking, as records were mostly outdated while North African habitats are undergoing fast changes (Samraoui et al., 2010). Over the last 15 years, additions to the Maghrebian odonate fauna have been accumulating steadily (Boudot, 2008; Juillerat & Monnerat, 2009; Korbaa, Ferreras-Romero, Bejaoui, & Boumaza, 2014; Samraoui & Alfarhan, 2015; Samraoui,

*Corresponding author. Email: bsamraoui@yahoo.fr

Weekers, & Dumont, 2003), culminating with the recent discovery of a species new to science in Morocco, *Onychogomphus boudoti* (Ferreira et al., 2014). This highlights the need for more surveys of the Maghrebian odonate fauna. This survey aims at improving our knowledge of the Gomphidae of Algeria and the Maghreb by (1) assessing their status, (2) identifying environmental factors driving their spatial and temporal distributions, and (3) documenting specific threats to their habitats.

Methods

Eight wadis were sampled across Algeria, stretching from Wadi Kébir-Est near the Tunisian border to Wadi Saoura, close to Morocco (Figure 1, Table 1). Six wadis (Seybouse, Sebaou, Isser, El Harrach, Tafna, and Saoura) were sampled monthly from April to October 2013. A series of 4–10 stations was sampled along the course of each wadi. Kébir-East (2013 and 2014) and Chioukh, Mount Edough (2014) were sampled sporadically. A series of 12 abiotic descriptors




Figure 1. Map of Algeria displaying the studied watershed and sampled localities; Locality codes in Table 1.

Table 1. GPS coordinates and altitude of localities where Gomphidae were recorded. Negative values are used for western longitudes. Record of *O. costae* at Kébir-Est dates from 1993 (Benchalel & Samraoui, 2012). Code is locality code from Figure 1.

Watershed	Locality	Code	Species	Latitude	Longitude	Alt. (m)
Mount Edough						
	Wadi Chioukh		<i>O. uncatus</i>	36° 55' 32.5"	7° 40' 58.9"	700
Kébir-Est	Kébir-Est		<i>P. genei</i> , <i>O. costae</i>	36° 46' 03"	8° 21' 52"	25
Seybouse	Chihani	S8	<i>G. lucasii</i> , <i>O. costae</i>	36° 39' 13"	7° 46' 57"	26
	Mellah		<i>O. costae</i>	36° 27' 32"	7° 42' 56"	87
	Halia	S7	<i>G. lucasii</i> , <i>O. costae</i>	36° 24' 49"	7° 36' 40"	144
	Nador	S6	<i>O. costae</i>	36° 27' 33"	7° 34' 18"	187
	Zimba		<i>O. f. unguiculatus</i>	36° 27' 49"	7° 29' 15"	205
	Héliopolis		<i>G. lucasii</i>	36° 30' 48"	7° 27' 02"	259
	El Fedjoudj	S5	<i>G. lucasii</i>	36° 28' 24"	7° 23' 07"	221
	Medjez Amar	S2	<i>G. lucasii</i> , <i>O. costae</i>	36° 26' 34"	7° 18' 39"	287
	Aïn Makhlouf	S1	<i>G. lucasii</i> , <i>O. costae</i>	36° 14' 27"	7° 18' 37"	599
	El Aar		<i>O. f. unguiculatus</i>	36° 13' 34"	7° 19' 11"	609
	Cheniour		<i>G. lucasii</i>	36° 14' 52"	7° 20' 36"	742
	Krab		<i>G. lucasii</i>	36° 07' 12"	7° 32' 47"	788
Isser						
	Wadi Djamaa	S1	<i>O. f. unguiculatus</i>	36° 26' 01"	3° 50' 50"	371
	Wadi Djamaa	S2	<i>O. f. unguiculatus</i>	36° 28' 08"	3° 44' 46"	328
	Wadi Isser	S3	<i>O. f. unguiculatus</i>	36° 31' 34"	3° 40' 45"	179
	Wadi Isser	S5	<i>O. f. unguiculatus</i>	36° 37' 21"	3° 34' 49"	72
Sebaou						
	Chaiïb	S4	<i>O. f. unguiculatus</i>	36° 42' 14"	4° 10' 35"	100
	Djemaa El Had	S7	<i>O. f. unguiculatus</i>	36° 35' 50"	4° 12' 04"	360
El Harrach						
	Magtaa Lazrag	S1	<i>O. f. unguiculatus</i>	36° 27' 53"	3° 00' 55"	220
	Hammam Melouène	S4	<i>O. f. unguiculatus</i>	36° 29' 22"	3° 03' 07"	158
	Bougara	S6	<i>O. f. unguiculatus</i>	36° 32' 06"	3° 03' 45"	92
	Baba Ali	S7	<i>O. f. unguiculatus</i>	36° 36' 50"	3° 04' 29"	29
	Baba Ali	S9	<i>O. f. unguiculatus</i>	36° 37' 33"	3° 03' 45"	25
Saoura						
	Djorf Torba	S2	<i>L. tetraphylla</i> , <i>O. costae</i>	31° 30' 34"	-2° 46' 25"	687
Tafna			No records of Gomphidae	34° 44' 13"	-1° 34' 57"	

were measured: water temperature (WatT in °C), air temperature (AirT in °C), dissolved oxygen (O2 in mg l⁻¹), per cent dissolved oxygen (O2per in %), pH, redox potential (Elec in mV), salinity (Sal in PSU), conductivity (Cond in mS cm⁻¹), speed (Speed in cm s⁻¹), bed width (Width in m), water depth (Depth in m) and altitude (Alt in m).

Adults and exuviae were recorded and identified using various keys (Dijkstra & Lewington, 2006; Gerken & Sternberg, 1999; Grand & Boudot, 2006). A principal component analysis (PCA) was performed on presence/absence data (21 stations/12 environmental variables and four species) using the package FactoMineR (Lê, Josse, & Husson, 2008). Data analyses and graphs were carried out using R (R Development Core Team, 2014).

Results

The present survey has yielded six species of Gomphidae (Figures 2, 3; Table 1). One species, the Critically Endangered *Lindenia tetraphylla*, recorded in the nineteenth century in eastern Algeria and deemed to be regionally extinct, was rediscovered in the western part of Algeria. Another species, tentatively assigned to *O. uncatus*, may represent either simply a pale phenotype of the latter or an undescribed taxon.




Figure 2. Map of Algeria with recorded localities of Gomphidae (inset).

Species data

1. *Gomphus lucasii* Selys, 1849

Status. VU.

Flight period. April–June.

Locality. Seybouse.

Previous localities. Rhummel, Oubeïra (Selys, 1849), Oran (Selys & Hagen, 1850, 1857), Orléansville (now Chlef) (Selys, 1871; Selys & Hagen, 1857), between Blida and Médéa (Kolbe, 1885), Constantine (McLachlan, 1897), Rhummel (Martin, 1901), Teniet el Had, Sebdou (Morton, 1905), Constantine, Oubeïra (Martin, 1910), Hammam Bou Hadjar (Schmidt, 1936), Numidia (Samraoui & Menaï, 1999).

Comments. This species is present from sea-level to foothills (600 m), but is particularly abundant at lower altitudes.

2. *Onychogomphus uncatus* (Charpentier, 1840)

Status. LC.

Flight period. July.

Locality. Wadi Chioukh (Séraïdi, Mount Edough).

Previous localities. Teniet el Had, Sebdou (Morton, 1905), Rhummel (Martin, 1910), Ighzer Temda (Lacroix, 1925), Tlemcen (Schmidt, 1936), Numidia, Mechroha (Samraoui & Menaï, 1999).

Comments. On 29 July 2014, two males of *O. uncatus* were recorded at Wadi Chioukh (Mount Edough) but only one of them could be collected. Both males were similar in having a pale thorax and differed enough from typical *O. uncatus* in their black/yellow pattern to warrant a



Figure 3. Photographs of (a) *Gomphus lucasii* (Wadi Seybouse, 2014); (b) *Lindenia tetraphylla* (Hijaz, Saudi Arabia, 2014); (c) *Onychogomphus forcipatus unguiculatus* (Wadi Seybouse, 2013); (d) immature *O. costae* (Wadi Seybouse, 2014).

brief description (Figure 4a, b) to stimulate further taxonomical investigations. The vertex is all-black, but the yellow “collar” is more narrowly interrupted in the middle by black. These males lack extensive confluent black lines on the side of the thorax, a feature which is reminiscent of *O. forcipatus unguiculatus* and, to a lesser extent, of *O. lefebvrei*. They differ in structure from the former by their “*O. uncatus*-type” epiproct, just semi-circular and missing any differentiated structure at the apex, and from the latter by the presence of a sub-basal tooth and a darker pterostigma. These males are clearly distinct from *O. boudoti*, recently described from Morocco (Ferreira et al., 2014) but more specimens are needed to confirm its distinctiveness from the usual European and Moroccan specimens of *O. uncatus*. So pending the collection of more specimens, the Edough taxon is assigned to *O. uncatus*, a species to which it is undoubtedly closely related. Males were settled on the ground on the shore of a mountain stream with a moderate flow. Habitat was shared mainly with patrolling *Boyeria irene*, *Aeshna cyanea* and *A. mixta* (Samraoui & Alfahhan, 2015).

3. *Onychogomphus forcipatus unguiculatus* (Vander Linden, 1820)

Status. LC.

Flight period. June–July.

Locality. Seybouse, El Harrach, Sebaou, Isser.

Previous localities. Algeria (Selys, 1849), between Médéa and Blida (Kolbe, 1885), Constantine (McLachlan, 1897), Sebdou (Morton, 1905), Constantine (Martin, 1910), Hammam Bou Hadjar (Schmidt, 1936), Numidia, Jijel (Samraoui & Menaï, 1999).



Figure 4. Side view of the Edough Onychogomphus assigned to *O. uncatus* (a). Dorsal view of the same specimen (b).

Comments. Relatively common in the centre of Algeria, but confined to a few sites in the east. At Wadi Seybouse, the species was more abundant at medium altitudes.

4. *Onychogomphus costae* Selys, 1885

Status. NT.

Flight period. May–October. Pairs in copula were noted in June and August.

Locality. Seybouse, Saoura.

Previous localities. Oran (Selys, 1871), Biskra, Constantine (McLachlan, 1897; Martin, 1901, 1910), Hammam Bou Hadjar (Schmidt, 1936), Bou Faadid, Reggan, Aoulef (Reymond, 1952), Numidia (Samraoui & Menai, 1999).

Comments. A species endemic to the Maghreb and the Iberian Peninsula. It is relatively common in northeast Algeria but appears to be less frequent further west. Our record, based on a single exuvia, confirms that the species may penetrate the Sahara. The old record of *Paragomphus costae* by Reymond (1952), ascribed to *P. genei* by Boudot et al. (2009), is thus upheld by this new finding. The same applies to the Tunisian record at Tozeur by Campion (1914), considered doubtful by Jödicke, Arlt, Kunz, Lopau, and Seidenbusch (2000).

5. *Paragomphus genei* (Selys, 1841)

Status. LC.

Flight period. October, but may be found from April to October (Samraoui & Menai, 1999).

Locality. Kébir-East watershed, Brabtia, close to Wadi Bouarroug.

Previous localities. Oubeïra, Biskra (McLachlan, 1897), Biskra (Martin, 1901, 1910; Ris, 1909–1913), Oubeïra (Martin, 1910), Algiers (Ander, 1929), Tassili n'Ajjer (Dumont, 1979), Numidia, Laghouat, El Goléa, Adrar, Timimoun (Samraoui & Menai, 1999).

Comments. Records under the synonyms of *O. hageni* Selys and *Mesogomphus hageni* Selys have been assigned to *P. genei*. Emergences seem to spread from May to November. A mass emergence was witnessed at Wadi Kébir-Est in September 1993, but the locality was heavily degraded in 2014 and the species was absent. An adult was recorded in a maquis near El Kala in October 2014.

6. *Lindenia tetraphylla* (Vander Linden, 1825)

Status. CR.

Flight period. May.

Locality. Saoura.

Previous localities. La Calle (Selys, 1849).

Comments. The species was first recorded in Algeria in the nineteenth century and was deemed extinct (Boudot et al., 2009). A population was discovered in nearby Tunisia (Kunz & Kunz, 2001). An exuvia was found downstream of the Djorf Torba reservoir, proof of the reproduction of the species in western Algeria.

Multivariate analysis

The range of occupied habitats by the recorded species exhibited significant variation of physicochemical variables (Figures 5, 6). Dimension reduction was performed using PCA. Four components, representing 76.4% of the total variance, were retained (Figure 7). The first component (29.7% of the inertia) is associated with high pH and large bed width. Habitats having these characteristics are linked to *G. lucasii*, *O. costae* and *L. tetraphylla* (Figure 7). This first component opposes habitats with high depth of water, occupied by *O. f. unguiculatus*. The second component (22.2% of the inertia) is associated with dissolved oxygen and temperature. The third component (14.2% of the inertia) opposes both *G. lucasii* and *O. costae* to *L. tetraphylla* on the basis of salinity and altitude (Figure 8). The fourth component (10.2% of the inertia) is of minor importance and separates sites on the basis of water current (Figure 8).

Discussion

Status

Six species, representing four genera (*Gomphus*, *Lindenia*, *Onychogomphus* and *Paragomphus*), are believed to exist in Algeria and they all have been recorded during this survey. A further two species are cited in the literature:

- (1) *Onychogomphus lefebvrei* (Rambur, 1842) was reported by Martin (1910) from Wadi Rhummel and Biskra. These records will be discussed in the light of the new finding of the pale form of *O. uncatus* from Mount Edough.
- (2) *Gomphus simillimus maroccanus* Lieftinck, 1966, of which the taxonomic status deserves further investigations (Dumont, 1972; Boudot et al., 2009), recorded in Algeria (Brauer, 1876 in Le Roi, 1915), at El Guerra and Wadi Seybouse (Martin, 1910), and at Tlemcen (Schmidt, 1936). The former three records are now considered to pertain to *G. lucasii* (Samraoui & Menai, 1999) but not the last one, as in the same paper dealing with




Figure 5. Extent of variation in bed width, depth of water and current speed of habitats occupied by *G. lucasii* (Gluc), *L. tetraphylla* (Ltet), *O. costae* (Ocost), and *O. forcipatus* (Ofor).




Figure 6. Extent of variation in altitude, water conductivity and pH of habitats occupied by *G. lucasii* (Gluc), *L. tetraphylla* (Ltet), *O. costae* (Ocost), and *O. forcipatus* (Ofor).




Figure 7. First two principal components of (a) variable factor map and (b) individual factor map. (c) Scree plot.

larvae/exuviae of Gomphidae, the author listed *G. lucasii* at Hammam Bou Hadjar and *G. simillimus (maroccanus)* at Tlemcen, and therefore discriminated the two taxa. However, the validity and distribution of *G. simillimus maroccanus* in the Maghreb remain an open question.

The newly described *O. boudoti*, a Critically Endangered species, is endemic to Morocco and is deemed to be facing a high risk of global extinction (Ferreira et al. 2014).




Figure 8. Third and fourth principal component of (a) variable factor map and (b) individual factor map.

The pale form of O. uncatus and the O. lefebvreii enigma

In North Africa and southern Spain, *O. uncatus*, not unlike *A. cyanea* (Ferrerías-Romero, Dolores Atienzar, & Corbet, 1999; Samraoui & Corbet, 2000a), is confined to mountain streams whereas in other parts of Europe, the species may occupy also lowland rivers and canals (Suhling, 1996). Its status in North Africa is listed as Least Concern on the IUCN North African Red List (Samraoui et al., 2010) but this evaluation rests on the abundance of the species in Morocco and the range believed to span Morocco, Algeria and Tunisia. Past records of *O. uncatus* from Algeria are old (Lacroix, 1925; Martin, 1910; Morton, 1905), misidentified (Khelifa et al., 2011) or based on larvae (Benchalel & Samraoui, 2012; Samraoui & Corbet, 2000a).

The presence of a pale form in Mount Edough points to the need to re-evaluate the taxonomic status of the genus in North Africa and to draw conservation consequences. We may speculate that early in the twentieth century, Gadeau de Kerville may have collected the pale form of *O. uncatus* (the Edough phenotype) in Tunisia which was then misidentified by Martin as *O. lefebvreii* (Gadeau de Kerville, 1908). This latter species is an Asiatic species ranging from the

Middle East to Afghanistan (Boudot et al., 2009) and it is unlikely that it ever existed in Algeria (Samraoui & Menai, 1999) or Tunisia (Jödicke et al., 2000) and even in Egypt, its so-called “type locality”. A few years later, Martin (1910) collected in Algeria the same pale phenotype of *O. uncatus*, which he again misidentified as *O. lefebvrei*. There is thus an urgent need to check Martin’s collection to evaluate these speculations.

Lindenia tetraphylla in North Africa

For decades, *L. tetraphylla* had been one of the least familiar odonates of the Palearctic region and was known only from a few specimens (Schorr, Schneider, & Dumont, 1998). New data are now accumulating (Boudot, 2014; Boudot et al., 2009; De Knijf, Vanappelghem, & Demolder, 2013; Kulijer, De Knijf, & Franković, 2013; Lopau, 2010; Wildermuth & Martens, 2014). The glaring gaps in our knowledge of its breeding habitats with a suggested dichotomy in the use of stagnant waterbodies and running waters in the west and in the east of the species’ range, respectively, are gradually being filled in (Popova, 1997). This nomadic species is well adapted to arid and semi-arid regions (Schneider, 1988; Schröter, 2011) as it breeds in waterbodies exhibiting a wide range of succession, salinity and temperature (Busse, 1993; Karaman, 1979; Krupp & Schneider, 1988). However, its larval habitat deserves further investigations (Boudot, 2014; Brochard & van der Ploeg, 2013).

The previous and only record from Algeria dates from the nineteenth century when Lucas collected three females near Lake Oubeïra, in northeast Algeria (Selys, 1849). With the old Spanish records from the Valencia province, the new Algerian record represents one of the most western point of the species’ range (Boudot et al., 2009) and may reflect a recent westward expansion in Algeria either due to climate change or, more likely, the creation of new habitats like dams and reservoirs. Due to the lack of field investigation in the Saoura valley since 1990 (Samraoui & Menai, 1999), it is impossible to know the date of settlement of this species in the locality considered. Considered as an Iranian element (Lohmann, 1981), *L. tetraphylla* has a similar distribution pattern as another Euroasiatic species, *Selysiothemis nigra* (Vander Linden, 1825), also found in arid and semi-arid regions (Schneider, 1981; Schorr et al., 1998). Both species have scattered populations in the western Mediterranean and both seem to be extending their range in North Africa (Boudot et al., 2009). However, this expansion is controversial as such apparent changes in ranges may be due to increasing field investigations. Also, their overall worldwide range does not appear to expand northwards significantly, at least in the west Mediterranean but perhaps also in southwestern Asia according to the paucity of ancient field investigations in parts of this area (Boudot, pers. comm.). In recent years, *L. tetraphylla* has frequently been recorded after an apparent decrease in various areas of southern Europe like the Balkans and Italy (Boudot, 2014; Brochard & van der Ploeg, 2013; Gashtarov & Beshkov, 2010; Hardersen & Leo, 2011; Stille, Stille, & Schröter, 2014; Utzeri, Belfiore, & Peels, 2006) and North Africa (Kunz & Kunz, 2001), with self-sustaining populations attested in several regions (e.g. Boudot et al., 2009). This recent increase in a near constant range is largely ascribed to an increase in available man-made habitats (e.g. Boudot, 2014; Brochard & van der Ploeg, 2013), an increase in field investigations (e.g. Skvortsov & Kuvaev, 2010) and an increase in standard scientific publications.

Ecology

Biological communities are structured by the environmental conditions of their habitats, described as habitat templets (Southwood, 1977). Thus, a good knowledge of the many physical and biological factors that may determine the abundance and spatial distribution of Odonata along the river continuum (Vannote, Minshall, Cummins, Sedell, & Cushing, 1980) is essential.

The larva is considered the most influential stage in the life cycle of Odonata (Corbet & Brooks, 2008; McPeck, 2008). Gomphid larvae are burrowers that ambush prey and evade predators while buried in the sediment (Corbet & Brooks, 2008; Suhling & Müller, 1996). An important ecological driver of gomphid larvae distribution is the substrate (Leipelt & Suhling, 2001; Suhling, 1994; 1996). However, the adult stage, through site selection and dispersal, may also shape population dynamics (Remsburg, 2011).

Although of a preliminary nature and constrained by a small sample size, our exploratory analysis identifies potential environmental factors that may separate Algerian Gomphidae. Three species (*O. uncatus*, *P. genei*, and *L. tetrphylla*) were only recorded once, but they clearly differed from the other gomphids on the basis of habitats. The former occupies mountainous streams whereas the latter two species, which are able to breed in coastal and inland standing waterbodies, are more frequent in arid and semi-arid regions. *Paragomphus genei* breeds in sandy parts of wadis with residual pools and sluggish current (Benchalel & Samraoui, 2012). A striking similarity is found in the habitat partitioning of *G. lucasii* and *O. forcipatus unguiculatus* in Algeria compared to that of *G. pulchellus* and *O. forcipatus unguiculatus* in southern Spain (Ferrerías-Romero & García-Rojas, 1995). In Spain, these two latter species are semivoltine and display some form of altitudinal separation by occupying different parts of the same river channel (Ferrerías-Romero & García-Rojas, 1995). Likewise, at Wadi Seybouse where they co-occur, *G. lucasii* is found more often in the lower reaches of the wadi and *O. forcipatus unguiculatus* is more frequent in the middle and upper parts. Another shared similarity between the two gomphid pairs across the Mediterranean is that they may share the same life histories. *Gomphus pulchellus* is a “spring species” (or type-1, *sensu* Corbet, 1999) in Spain, like *G. lucasii* in Algeria, whereas *O. forcipatus unguiculatus* behaves as a “summer species” (or type-2, *sensu* Corbet, 1999) in both countries (Ferrerías-Romero & García-Rojas, 1995; this study). Thus, there is a remarkable convergence in the life histories of *G. lucasii* and *G. pulchellus*, which may be regarded as vicariant species.

Spatial partitioning is not the only mean of coexistence in local gomphids. *Gomphus lucasii* and *O. costae*, which often share the same habitats, have distinct flight periods. The very long flight period of *O. costae* is reminiscent of that of *Anax imperator* in Britain (Corbet, 1957), which exhibits splitting larval cohorts: semivoltine individuals emerging early in the season as in “spring species” and univoltine individuals with fast development emerging later in the season as in “summer species”. The suggested life history pattern of *O. costae* in Algeria is supported by studies in southern Spain (Cano-Villegas & Ferrerías-Romero, 2005). This temporal partitioning which may ease competition is exhibited by many congeneric Mediterranean odonates (Ferrerías-Romero & García-Rojas, 1995; Samraoui, 2009; Samraoui & Corbet, 2000b).

Conservation

The alarming status of freshwater species in North Africa (Riservato et al., 2009; Samraoui et al., 2010) is intimately associated with explosive demography and urbanization (Tabutin, Vilquin, & Biraben, 2002), combined with the increasing water demands of agriculture, industry development and domestic use (Benoit & Comeau, 2005). The lack of sewage treatment combined with human encroachment has led to pollution reaching a critical point. The proliferation of dam construction to meet the heavy demand for water and the lack of law enforcement against gravel extraction in rivers bed has worsened the water quality of most wadis in North Africa and the Middle East (Jödicke, Boudot, Jacquemin, Samraoui, & Schneider, 2004; Schneider & Schneider, 2010). Global warming may exacerbate the situation but may prompt northern African countries to take steps towards a rational management of water resources.

This survey is of a preliminary nature and further investigations, involving a greater array of wadis, are required. Combined sampling of adults and exuviae proved worthwhile in the present

study as Gomphidae are strong flyers and adults may easily evade detection (Brochard & van der Ploeg, 2013; Raebel, Merckx, Riordan, Macdonald, & Thompson, 2010). We clearly need more surveys and systematic monitoring to strengthen our knowledge of the status of Maghrebian Odonata and improve our understanding of the processes that regulate their abundance and distribution. We also need to enforce prompt and effective conservation measures to avert threats that imperil their survival (Clausnitzer et al., 2012).

Acknowledgements

We are most grateful to J.-P. Boudot, M. Ferreras-Romero and H. Dumont for making helpful suggestions which greatly improved our manuscript. We also thank two anonymous referees for instructive comments. The research was funded by the National Plan for Science and Technology, Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia under project grant no. 12-ENV2569-02.

ORCID

Boudjéma Samraoui  <http://orcid.org/0000-0002-0608-9021>

References

- Ander, K. E. V. (1929). Über die Nympe von *Mesogomphus Hageni* Selys. *Konowia* (Vienna), 8, 159–162. Retrieved from <http://www.landesmuseum.at/biophp/de/konowia.pdf>
- Benchalel W., & Samraoui B. (2012). Caractérisation écologique et biologique de l'odonatofaune de deux cours d'eau méditerranéens : l'oued El-Kébir et l'oued Bouaroug (Nord-Est de l'Algérie). *Méditerranée*, 118, 19–27. <http://dx.doi.org/10.4000/mediterranee.6182>
- Benoit, G., & Comeau, A. (2005). *A sustainable future for the Mediterranean: the Blue Plan's environment and development outlook*. Plan Bleu: Earthscan. Retrieved from <http://dx.doi.org/10.4324/9781849770323>
- Boudot, J.-P. (2008). *Selysiothemis nigra* (Vander Linden, 1825), nouveau pour le Maroc, et autres observations sur les Odonates du Maghreb nord-occidental (Odonata: Anisoptera : Libellulidae). *Martinia*, 24, 3–29.
- Boudot, J.-P. (2014). A brief observation of egg laying in *Lindenia tetraphylla* (Odonata: Gomphidae) on Kríti (Crete), Greece. *Notulae Odonatologicae*, 8, 94–96.
- Boudot, J.-P., Kalkman, V. J., Azpilicueta Amorin, M., Bogdanovic, T., Cordero Rivera, A., Degabriele, G., ... Schneider, W. (2009). Atlas of the Odonata of the Mediterranean and North Africa. *Libellula Supplement*, 9, 1–256.
- Brochard, C., & van der Ploeg, E. (2013). Searching for exuviae of endemic Odonata species in Greece. *Brachytron*, 15, 83–101.
- Busse, R. (1993). Libellen von der türkischen Südküste. *Libellula*, 12, 39–46.
- Campion, H. (1914). List of Odonata collected at Tozeur, South Tunisia, by Mr G C Champion in May, 1913. *Entomologist's Monthly Magazine*, 2, 118.
- Cano-Villegas, F. J., & Ferreras-Romero, M. (2005). *Contribution to knowledge of the biology of Onychogomphus costae Selys, 1885 (Odonata: Gomphidae) in southern Spain*. Poster presentation. Abstracts Book 4th WDA International Symposium of Odonatology, Pontevedra, 26–30 July 2015.
- Clausnitzer, V., Dijkstra, K.-D. B., Koch, R., Boudot, J.-P., Darwall, W. R. T., Kipping, J., ... Suhling, F. (2012). Focus on African freshwaters: hotspots of dragonfly diversity and conservation concerns. *Frontiers in Ecology and Environment*, 10, 129–134. <http://dx.doi.org/10.1890/110247>
- Corbet, P. S. (1957). The life history of the Emperor Dragonfly *Anax imperator* Leach (Odonata: Aeshnidae). *Journal of Animal Ecology*, 26, 1–69. <http://dx.doi.org/10.2307/1781>
- Corbet, P. S. (1999). *Dragonflies: Behavior and ecology of Odonata*. Ithaca: Cornell University Press. <http://dx.doi.org/10.1076/aqin.23.1.83.4929>
- Corbet, P. S., & Brooks, S. (2008). *Dragonflies*. London: Collins.
- De Knijf, G., Vanappelpghem, C., & Demolder, H. (2013). Odonata from Montenegro, with notes on taxonomy, regional diversity and conservation. *Odonatologica*, 42, 1–29.
- Dijkstra, K.-D. B., & Lewington, R. (2006). *Field guide to the dragonflies of Britain and Europe*. Gillingham: British Wildlife Publishing.
- Dumont, H. J. (1972). Contribution à la connaissance des odonates du Maroc. *Bulletin de la Société des Sciences Naturelles et Physiques du Maroc*, 52, 149–179.
- Dumont, H. J. (1979). *Limnologie van Sahara en Sahel*. Gent: Rijksuniversiteit.
- Ferreira, S., Velo-Anton, G., Brochard, C., Vieira, C., Celio Alves P., Thompson, D. J., ... Brito, J. C. (2014). A critically endangered new dragonfly species from Morocco: *Onychogomphus boudoti* sp. nov. (Odonata: Gomphidae). *Zootaxa*, 3856, 349–365. <http://dx.doi.org/10.11646/zootaxa.3856.3.3>

- Ferreras-Romero, M., Dolores Atienzar, M., & Corbet, P. S. (1999). The life cycle of *Onychogomphus uncatus* (Charpentier, 1840) (Odonata: Gomphidae) in the Sierra Morena Mountains (southern Spain): An example of protracted larval development in the Mediterranean basin. *Archiv für Hydrobiologie*, 144, 215–228.
- Ferreras-Romero, M., & García-Rojas, A. M. (1995). Life-history patterns and spatial separation exhibited by the odonates from a Mediterranean inland catchment in southern Spain. *Vie et Milieu*, 45, 157–166.
- Ferreras-Romero, M., Márquez-Rodríguez, J., & Ruiz-García, A. (2009). Implications of anthropogenic disturbance factors on the Odonata assemblage in a Mediterranean fluvial system. *International Journal of Odonatology*, 12, 413–428. <http://dx.doi.org/10.1080/13887890.2009.9748354>
- Gadeau de Kerville, H. (1908). *Odonates. Voyage Zoologique en Khroumirie (Tunisie). Mai-Juin, 1906*. Odonata, 59–61. Paris: Baillière.
- Gashtarov V., & Beshkov S. (2010). *Lindenia tetraphylla* (Vander Linden, 1825) (Odonata: Gomphidae) a new genus and species for the Bulgarian fauna. *The Entomologist's Record and Journal of Variation*, 122, 272–274.
- Gerken, B., & Sternberg, K. (1999). Die Exuvien Europäischer Libellen (Insecta, Odonata). Höxter: Verlag und Werbeagentur. http://dx.doi.org/10.1007/978-3-8274-2452-5_13
- Grand, D., & Boudot J.-P. (2006). *Les libellules de France, Belgique et Luxembourg*. Mèze: Biotope.
- Hardersen, S., & Leo, P. (2011). Dragonflies of Iglesias (SW Sardinia) and additional records of rare or poorly known species from Sardinia (Odonata). *Conservazione Habitat invertebrate*, 5, 243–253.
- Jödicke, R., Arlt, J., Kunz, B., Lopau, W., & Seidenbusch, R. (2000). The Odonata of Tunisia. *International Journal of Odonatology*, 3, 41–71. <http://dx.doi.org/10.1080/13887890.2000.9748135>
- Jödicke, R., Boudot, J. P., Jacquemin, G., Samraoui, B., & Schneider, W. (2004). Critical species of Odonata in northern Africa and the Arabian Peninsula. *International Journal of Odonatology*, 7(2), 239–253. <http://dx.doi.org/10.1080/13887890.2004.9748213>
- Juillerat, L., & Monnerat, C. 2009. Odonata in southern Morocco, with first records of *Orthetrum ransonnetii* and *Sympetrum sinaiticum* (Odonata: Libellulidae). *Libellula*, 28, 97–115.
- Karaman, B. S. (1979). Contribution à la connaissance de l'écologie des larves d'odonates dans l'écosystème du lac de Doiran. *Annales de la Faculté de Biologie de l'Université de Skopje*, 32, 191–199.
- Khelifa, R., Youcefi, A., Kahleras, A., Alfarhan, A. H., Al-Rasheid, K. A. S., & Samraoui, B. (2011). L'odonatofaune de la Seybouse en Algérie: intérêt pour la biodiversité du Maghreb (Insecta: Odonata). *Revue d'Ecologie (Terre et Vie)*, 66, 55–66.
- Kolbe, H. J. (1885). Beitrag zur Kenntniss der Pseudoneuroptera Algeriens und der Ostpyrenäen. *Berliner entomologische Zeitschrift*, 29, 151–157. <http://dx.doi.org/10.1002/mmnd.18850290126>
- Korbua, M., Ferreras-Romero, M., Bejaoui, M., & Boumaza, M. (2014). Two species of Odonata newly recorded from Tunisia. *African Entomology*, 22, 291–296. <http://dx.doi.org/10.4001/003.022.0212>
- Krupp, F., & Schneider, W. (1988). Die Süßwasserfauna des Vorderen Orients. Anpassungs- strategien und Besiedlungsgeschichte einer zoogeographischen Übergangszone. *Natur und Museum*, 118, 193–213.
- Kulijer, D., De Knijf, G., & Franković, M. (2013). Review of the Odonata of Bosnia and Herzegovina. *Odonatologica*, 42, 109–123.
- Kunz, B., & Kunz, D. (2001). *Lindenia tetraphylla* Wiederfund für Nordafrika (Odonata: Gomphidae). *Libellula*, 20, 79–85.
- Lacroix, J. L. (1925). Quelques Névroptères (sens. Lat.) d'Afrique. *Bulletin de la Société d'Histoire naturelle d'Afrique du Nord*, 16, 258–263.
- Lê, S., Josse, J., & Husson, F. (2008). FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software*, 25, 1–18.
- Leipelt, K. G., & Suhling, F. (2001). Habitat selection of larval *Gomphus graslinii* and *Oxygastra curtisii* (Odonata: Gomphidae, Corduliidae). *International Journal of Odonatology*, 4, 23–34. <http://dx.doi.org/10.1080/13887890.2001.9748155>
- Le Roi, O. (1915). Odonaten aus der Algerischen Sahara von der Reise von Freiherrn H. Geyr von Schweppenburg. Mit einer Übersicht der Nordafrikanischen Odonaten-Fauna. *Deutsch Entomologische Zeitschrift*, 1915, 609–634. <http://dx.doi.org/10.1002/mmnd.48019150606>
- Lohmann, H. (1981). Postglaziale Disjunktionen bei europäischen Libellen. *Libellula*, 1, 2–4.
- Lopau, W. (2010). Verbreitungsatlas der Libellen in Griechenland (Odonata). *Libellula Supplement*, 10, 5–153.
- Martin, R. (1901). Les odonates en Algérie au mois de mai. *Feuille des Jeunes Naturalistes, Paris*, 31, 249–250.
- Martin, R. (1910). Contribution à l'étude des Neuroptères de l'Afrique. II. Les odonates du département de Constantine. *Annales de la Société Entomologique de France*, 79, 82–104.
- McLachlan, R. (1897). Odonata collected by the Rev. E.A. Eaton in Algeria, with annotations. *Entomologist's Monthly Magazine*, S.2, 8, 152–157.
- McPeck, M. A. (2008). Ecological factors limiting the distributions and abundances of Odonata. In A. Córdoba-Aguilar (Ed.), *Dragonflies and damselflies. Model organisms for ecological and evolutionary research*. Oxford: Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780199230693.003.0005>
- Morton, K. J. (1905). Odonata collected by Miss M. Fountaine in Algeria, with description of a new species of *Ischnura*. *Entomology Monthly Magazine*, S.2, 16, 146–149.
- Popova, O. N. (1997). *Dragonflies of the Caucasus*. Abstracts of Papers and Posters presented at the 14th International Symposium of Odonatology 1997, Maribor, p. 31.
- R Development Core Team (2014). *R: A language and environment for statistical computing*. Vienna, Austria: The R Foundation for Statistical Computing.

- Raebel, E. M., Merckx, T., Riordan, P., Macdonald, D. W., & Thompson, D. J. (2010). The dragonfly delusion: why it is essential to sample exuviae to avoid biased surveys. *Journal of Insect Conservation*, 14, 523–533. <http://dx.doi.org/10.1007/s10841-010-9281-7>
- Remsburg, A. (2011). Relative influence of prior life stage and habitat variables on dragonflies (Odonata: Gomphidae) densities among lake sites. *Diversity*, 3, 200–216. <http://dx.doi.org/10.3390/d3020200>
- Reymond, A. (1952). Insectes de divers ordres récoltés au Sahara central au cours d'une mission du Centre national de la Recherche scientifique en 1947–1948. *Bulletin de la Société Sciences Naturelles du Maroc*, 32, 77–89.
- Ris, F. (1909–1913). *Catalogue systématique et descriptif des Collections Zoologiques du Baron Edm. de Selys Longchamps*. Bruxelles: Libellulinen.
- Riservato, E., Boudot, J.-P., Ferreira, S., Jović, M., Kalkman, V. J., Schneider, W., Samraoui, B., & Cuttelod, A. (2009). *The status and distribution of dragonflies of the Mediterranean Basin*. Gland: IUCN. <http://dx.doi.org/10.2305/iucn.ch.2009.18.en>
- Samraoui, B. (2009). Seasonal ecology of Algerian Lestidae (Odonata). *International Journal of Odonatology*, 12, 383–394. <http://dx.doi.org/10.1080/13887890.2009.9748352>
- Samraoui, B., & Alfathan, A. H. (2015). Odonata of mountain streams from Mount Edough, Algeria, and Kroumiria, Tunisia. *African Entomology*, 23, 172–179. <http://dx.doi.org/10.4001/003.023.0129>
- Samraoui, B., Boudot, J.-P., Ferreira, S., Riservato, E., Jović, M., Kalkman, V. J., & Schneider, W. (2010). The status and distribution of dragonflies. In N. Garcia, A. Cuttelod & D. Abdul Malak (Eds), *The status and distribution of freshwater biodiversity in Northern Africa* (pp. 51–70). Gland, Switzerland: IUCN. Retrieved from http://cmsdata.iucn.org/downloads/the_status_and_distribution_of_freshwater_biodiversity_in_northern_africa.pdf
- Samraoui, B., & Corbet, P. S. (2000a). The Odonata of Numidia. Part I: Status and distribution. *International Journal of Odonatology*, 3, 11–25. <http://dx.doi.org/10.1080/13887890.2000.9748133>
- Samraoui, B., & Corbet, P. S. (2000b). The Odonata of Numidia. Part II: Seasonal ecology. *International Journal of Odonatology*, 3, 27–39. <http://dx.doi.org/10.1080/13887890.2000.9748134>
- Samraoui, B., & Menai, R. (1999). A contribution to the study of Algerian Odonata. *International Journal of Odonatology*, 2, 145–165. <http://dx.doi.org/10.1080/13887890.1999.9748126>
- Samraoui, B., Weekers, P. H. H., & Dumont, H. J. (2003). Two taxa within the North African *Lestes virens* complex. *Odonatologica*, 32, 131–142.
- Schmidt, E. (1936). Die Westpaläarktischen Gomphiden-Larven nach ihren letzten Häufen (Ins. Odon.). *Senckenbergiana*, 18, 270–282.
- Schneider, W. (1981). Eine Massenwanderung von *Selysiothemis nigra* (Vander Linden, 1825) (Odonata: Macrodiplacidae) und *Lindenia tetraphylla* (Vander Linden, 1825) (Odonata: Gomphidae) in SüdJordanien. *Entomologische Zeitschrift*, 91, 97–102.
- Schneider, W. (1988). Dragonflies (Odonata) of the Wahiba Sands and adjacent areas, Eastern Oman. *Journal of Oman Studies, Special Report*, 3, 377–288.
- Schneider, T., & Schneider, J. (2010). Occurrence, behavior, and habitat preference of the Levant Pincertail, *Onychogomphus macrodon* Selys, 1887 in Turkey. *Zoology in the Middle East*, 49, 79–88. <http://dx.doi.org/10.1080/09397140.2010.10638393>
- Schorr, M., Schneider, W., & Dumont, H. J. (1998). Ecology and distribution of *Lindenia tetraphylla* (Insecta, Odonata, Gomphidae): A review. *International Journal of Odonatology*, 1, 65–88. <http://dx.doi.org/10.1080/13887890.1998.9748093>
- Schröter, A. (2011). A mass migration of *Aeshna affinis* in southern Kyrgyzstan: attempt to provide a spatial and temporal reconstruction (Odonata: Aeshnidae). *Libellula*, 30, 203–232.
- Selys-Longchamps, E. de (1849). Libellulinae. In P. H. Lucas (Ed.), *Exploration scientifique de l'Algérie. Zoologie. 2. Histoire naturelle des animaux articulés. Part 3, Insectes* (pp. 110–140). Paris: Imprimerie Nationale.
- Selys-Longchamps, E. de (1871). Nouvelle révision des odonates de l'Algérie. *Annales de la Société entomologique belge*, 14, 9–20.
- Selys-Longchamps, E. de, & Hagen, H. A. (1850). Revue des Odonates ou Libellules d'Europe. *Mémoires de la Société Royale des Sciences de Liège*, 6, 1–408.
- Selys-Longchamps, E. de, & Hagen, H. A. (1857). Monographie des Gomphines. *Mémoires de la Société Royale des Sciences de Liège*, 11, 257–720.
- Skvortsov, V. E., & Kuvaev, V. (2010). *Ischnura fountaineae* Morton, *Lindenia tetraphylla* (Vander Linden) and *Selysiothemis nigra* (Vander Linden) new for European Russia (Zygoptera: Coenagrionidae; Anisoptera: Gomphidae, Libellulidae). *Notulae odonatologicae*, 7, 49–51.
- Southwood, T. R. E. (1977). Habitat, the template for ecological strategies? *Journal of Animal Ecology*, 46, 337–365. <http://dx.doi.org/10.2307/3817>
- Stille, M., Stille, B., & Schröter, A. (2014). *Lindenia tetraphylla* – new for the island of Kérkira (Corfu), Greece (Odonata: Gomphidae). *Notulae odonatologicae*, 8, 77–116.
- Suhling, F. (1994). Spatial distribution of the larvae of *Gomphus pulchellus* Selys (Anisoptera: Gomphidae). *Advances in Odonatology*, 6, 101–111.
- Suhling, F. (1996). Interspecific competition and habitat selection by the riverine *Onychogomphus uncatus*. *Freshwater Biology*, 35, 209–217. <http://dx.doi.org/10.1046/j.1365-2427.1996.00491.x>
- Suhling, F., & Müller, O. (1996). *Die Flussjungfern Europas*. Heidelberg: Spektrum Akademischer Verlag.

- Tabutin, D., Vilquin, E., & Biraben, J.-N. (2002). *L'histoire de la population de l'Afrique du Nord pendant le deuxième millénaire*. Université catholique de Louvain, Département des sciences de la population et du développement, Document de travail, 15. Louvain-la-Neuve.
- Utzeri, C., Belfiore, C., & Peels, F. (2006). Some new records of *Lindenia tetraphylla* (Vander Linden) in Italy (Anisoptera: Gomphidae). *Notulae Odonatologicae*, 6, 90–92.
- Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., & Cushing, C. E. (1980). The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37, 130–137. <http://dx.doi.org/10.1139/f80-017>
- Wildermuth, H., & Martens, A. (2014). *Taschenlexikon der Libellen Europas*. Wiebelsheim: Quelle & Meyer.
- Woodward, G., & Hildrew, A. G. (2001). Invasion of a stream food web by a new top predator. *Journal of Animal Ecology*, 70, 273–288. <http://dx.doi.org/10.1111/j.1365-2656.2001.00497.x>
- Woodward, G., & Hildrew, A. G. (2002). The impact of a sit-and-wait predator: separating consumption and prey emigration. *Oikos*, 99, 409–418. <http://dx.doi.org/10.1034/j.1600-0706.2002.11210.x>